



## Class Problem – Astronomical Observation and Instrumentation



**Fig. 5.** Combined EPIC MOS spectra and best-fit models for sources X11. The spectra of X11 are fitted with one temperature thermal plasma model. The fit residuals for all sources are displayed in the bottom panels of each plot, with error bars representing  $1\sigma$  uncertainties.

**Figure 6.** (Left) spectrum (blue) and fit (red) of SN 1986J, ObsID 794. Lines of Mg and Si are visible in the fit. (Right) The XMM PN spectrum (blue) and fit (red) of SN 2005kd, XMM ID 0410581101. In this case, the Fe K line at around 6.7 keV, and a Ca line at  $\approx 4$  keV, are both apparent. The fit is consistent with that in [13]. In both of these Type IIs, the spectra are clearly thermal.



Sources:

<https://arxiv.org/pdf/2506.09120> and <https://www.mdpi.com/2218-1997/11/5/161>

- What portion of the EM spectrum is being studied in Figures 5 and 6 above? (might block out captions for this)
- Using the table, identify the elements seen in the three objects.

element	Energy (Kev)	element	Energy (Kev)	element	Energy (Kev)
O	0.18	Mg	1.33	Ar	3.32
Mg	0.25	Mg	1.45	Ar	3.69
Mg	0.27	Fe	1.66	Ca	3.86
O	0.64	Si	1.87	Ca	3.89
O	0.66	Si	1.98	Ca	4.11
Fe	0.80	Si	2.14	Ca	4.95
Fe	0.81	S	2.42	Fe	6.47
Ne	0.92	S	2.44	Fe	6.54
Ne	0.93	S	2.63	Fe	6.97
Ne	1.02	Ar	3.10	Fe	7.80

- Based on these elements, what type of supernova did each of these remnants come from? Justify your answer.
- If the spectra of X11 is assumed to be completely thermal, what temperature would its source be?
- If this same temperature is instead caused by shocks within the Snc remnant, what would be the material's initial velocity if the material is stationary after the shock? (Assume a pure Hydrogen material,  $m_H = 1.67 \times 10^{-27}$  kg and  $1\text{eV} = 1.602 \times 10^{-19}$  J)
- Both sets of data were taken with Chandra, can you draw a rudimentary diagram of the telescope?
- Finally, if the detector collects a field of view  $\pm 0.5$  deg but is the physical size of  $16.9' \times 16.9'$ , what total deflection does Chandra's most external mirrors (aperture of 1.2 m) need to apply to incoming light if the optical bench is 10 m?